

Basic = Intermediate

Baseline risk assessment

<p>Age:</p> <input type="text"/> <input type="text"/> yr.	<p>Pre-enrollment secondary insults:</p> <p>Hypoxia <input type="radio"/> No <input type="radio"/> Yes</p> <p>Hypotension <input type="radio"/> No <input type="radio"/> Yes</p>	<p>Pupillary reactivity:</p> <p><input type="radio"/> Both pupils reactive</p> <p><input type="radio"/> One non-reacting pupil</p> <p><input type="radio"/> Both pupils non-reactive</p>	<p>CT parameters:</p> <p>CT class <input type="text"/> (1-5)</p> <p>tSAH <input type="radio"/> No <input type="radio"/> Yes</p>
<p>Qualifying Motor Score For study admission: <input type="text"/> (1-6)</p>	<p>Time of Assessment:</p> <p><input type="radio"/> Post-stabilization</p> <p><input type="radio"/> Admission</p> <p><input type="radio"/> First hospital</p> <p><input type="radio"/> Scene of accident</p> <p><input type="radio"/> Other</p>	<p>Conditions of Assessment:</p> <p><input type="radio"/> No sedation/paralysis</p> <p><input type="radio"/> After stopping sedation</p> <p><input type="radio"/> After pharmacologic reversal</p> <p><input type="radio"/> Under sedation</p>	

Baseline risk assessment

Age: <input type="text"/> <input type="text"/> yr.	Pre-enrollment secondary insults: Hypoxia <input type="radio"/> No <input type="radio"/> Yes Hypotension <input type="radio"/> No <input type="radio"/> Yes	Pupillary reactivity: <input type="radio"/> Both pupils reactive <input type="radio"/> One non-reacting pupil <input type="radio"/> Both pupils non-reactive	CT parameters: CT class <input type="text"/> (1-5) tSAH <input type="radio"/> No <input type="radio"/> Yes			
<table><tr><td data-bbox="43 412 510 712">Qualifying Motor Score For study admission: <input type="text"/> (1-6)</td><td data-bbox="510 412 833 712">Time of Assessment: <input type="radio"/> Post-stabilization <input type="radio"/> Admission <input type="radio"/> First hospital <input type="radio"/> Scene of accident <input type="radio"/> Other</td><td data-bbox="833 412 1387 712">Conditions of Assessment: <input type="radio"/> No sedation/paralysis <input type="radio"/> After stopping sedation <input type="radio"/> After pharmacologic reversal <input type="radio"/> Under sedation</td></tr></table>				Qualifying Motor Score For study admission: <input type="text"/> (1-6)	Time of Assessment: <input type="radio"/> Post-stabilization <input type="radio"/> Admission <input type="radio"/> First hospital <input type="radio"/> Scene of accident <input type="radio"/> Other	Conditions of Assessment: <input type="radio"/> No sedation/paralysis <input type="radio"/> After stopping sedation <input type="radio"/> After pharmacologic reversal <input type="radio"/> Under sedation
Qualifying Motor Score For study admission: <input type="text"/> (1-6)	Time of Assessment: <input type="radio"/> Post-stabilization <input type="radio"/> Admission <input type="radio"/> First hospital <input type="radio"/> Scene of accident <input type="radio"/> Other	Conditions of Assessment: <input type="radio"/> No sedation/paralysis <input type="radio"/> After stopping sedation <input type="radio"/> After pharmacologic reversal <input type="radio"/> Under sedation				
Laboratory values: <input type="radio"/> Hemoglobine: <input type="text"/> <input type="text"/> · <input type="text"/> <input type="radio"/> mMol/L <input type="radio"/> mg/100 ml <input type="radio"/> Glucose: <input type="text"/> <input type="text"/> · <input type="text"/> <input type="radio"/> mMol/L <input type="radio"/> mg/100 ml <input type="radio"/> Prothrombine time: <input type="text"/> <input type="text"/> · <input type="text"/> sec.						